

December 22, 2006

JINR E2–2006–166

Disorder parameter and canonical quantization approach in lattice gauge theory

V.K. Mitrjushkin^{a,b}

^a *Joint Inst. for Nuclear Research, 141980 Dubna, Russia*

^b *Inst. of Theor. and Exper. Physics, 117259 Moscow, Russia*

Abstract

This work is dedicated to the derivation of the 'functional' integral representation of the disorder parameter $\langle \hat{\mu} \rangle$, $\hat{\mu}$ being the disorder *operator*. This derivation resolves the problem of the choice of the so called modified action as well as the problem of the choice of the boundary conditions along the 'time' direction. The question of the gauge fixing in the functional integral representation is also discussed.

1 Introduction

The concept of the disorder parameter [1] looks very promising in lattice gauge theories. In particular, a properly defined disorder parameter $\langle \hat{\mu} \rangle$ gives a possibility to check if the dual superconductivity of the ground state and monopole condensation are responsible for the color confinement [2]–[4]. Last years a number of papers has been dedicated to the (numerical) study of this hypothesis (see, e.g, papers [5] – [14] and also papers [15, 16] where the expectation value of the monopole creation operator of the Fröhlich-Marchetti type [17] was shown to be a disorder parameter as well).

Very often (see, e.g., [6]), the disorder parameter $\langle \hat{\mu} \rangle$ is defined as an average of a disorder *operator* $\hat{\mu}$:

$$\langle \hat{\mu} \rangle \sim \text{Tr} (\dots \times \hat{\mu}) \ ;$$

(for exact definitions see below). On the other hand, in the functional integral representation (Wilson's approach) the disorder parameter is written as

$$\langle \hat{\mu} \rangle = Z^{-1} \int [dU] \tilde{\mu}(U) \cdot \exp \left\{ -S_W(U) \right\} \equiv Z^{-1} \int [dU] \exp \{ -\tilde{S}(U) \} \ , \quad (1.1)$$

where $S_W(U)$ is the standard Wilson action and the partition function is given by $Z = \int [dU] \cdot \exp\{-S_W(U)\}$.

Therefore, the first question of interest is the connection between operator $\hat{\mu}$ and corresponding functional $\tilde{\mu}(U)$ (or $\tilde{S}(U)$).

Another two questions of interest are the following.

- What kind of boundary conditions along the forth (i.e. imaginary time) direction should/could one use in the functional integral representation (note that operator $\hat{\mu}$ is defined in the $3d$ space while functional integral is defined in the $4d$ space) ?
- Should one fix any gauge in the functional integral ?

The next section contains the brief discussion of the canonical quantization approach on the lattice and the transfer matrix formalism. The third section is dedicated to the disorder parameter. The last section is reserved for conclusions.

In what follows the periodic boundary conditions in the spacelike directions are presumed. The gauge groups is $SU(N)$, symbol a_4 means spacing in the forth direction.

2 Canonical quantization and transfer matrix

Hamiltonian acts in the space $\mathcal{H} = \{|\Psi\rangle\}$ where $\langle\Psi|\Psi\rangle < \infty$. The wave function of the state $|\Psi\rangle$ depends on link variables U_{l_s} where l_s are spacelike links ($l_s \equiv (\vec{x}; \vec{y}) \equiv (\vec{x}; \vec{x} + \hat{k})$). Therefore, $|\Psi\rangle$ can be represented in the form

$$|\Psi\rangle = \int \prod_{l_s} dU_{l_s} \Psi(U) |U\rangle, \quad (2.1)$$

where $|U\rangle$ is the eigenstate of the operators $\hat{U}_{l_s} = \hat{U}_{\vec{x};\vec{y}} : \hat{U}_{l_s}|U\rangle = U_{l_s}|U\rangle$, and

$$\langle U'|U\rangle = \prod_{l_s} \delta(U'_{l_s}; U_{l_s}); \quad \int [dU] |U\rangle \langle U| = 1. \quad (2.2)$$

Let $\Omega = \exp\{i \sum_a \omega^a T^a\} \in SU(N)$ be some matrix and for every spacelike link l let us define unitary operators $R_l(\Omega) : R_l(\Omega)|U\rangle = |U'\rangle$; $R_l(\hat{1}) = \hat{1}$, where $U'_l = \Omega U_l$ and $U'_{l_1} = U_{l_1}$ for $l_1 \neq l$. It is easy to show,

$$R_l(\Omega) \hat{U}_l R_l^\dagger(\Omega) = \Omega^\dagger \hat{U}_l. \quad (2.3)$$

Operators $R_l(\Omega)$ can be represented in the form $R_l(\Omega) = \exp\{i \sum_a \omega_l^a E_l^a\}$, where hermitian operators E_l^a have the following commutation relations

$$[E^a, E^b] = i f^{abc} E^c; \quad [E^a, \hat{U}] = -T^a \hat{U}; \quad [E^a, \hat{U}^\dagger] = \hat{U}^\dagger T^a. \quad (2.4)$$

One can establish a connection between $E_l^a = E_{\vec{x};\vec{y}}^a$ and $E_{-l}^a = E_{\vec{x}+\hat{j};-\hat{j}}^a = E_{\vec{y};\vec{x}}^a$:

$$E_{-l}^a = - \sum_b G^{ab}(U_l) \cdot E_l^b ; \quad G^{ab}(U_l) = 2\text{Tr}(U_l T^a U_l^\dagger T^b) , \quad (2.5)$$

Evidently,

$$[E_{-l}^a, E_{-l}^b] = -if^{abc} E_{-l}^c ; \quad [E_{-l}^a, \hat{U}_l] = \hat{U}_l T^a ; \quad [E_{-l}^a, \hat{U}_l^\dagger] = -T^a \hat{U}_l^\dagger . \quad (2.6)$$

The generator $\Gamma^a(\vec{x})$ of the gauge transformation in $\vec{x}$ and the operator $\mathcal{R}(\Omega)$ of the gauge transformation are given by

$$\Gamma^a(\vec{x}) \equiv \sum_{k=1}^3 \left(E_k^a(\vec{x}) + E_{-k}^a(\vec{x}) \right) ; \quad \mathcal{R}(\Omega) = \prod_{\vec{x}} e^{i \sum_a \omega^a(\vec{x}) \Gamma^a(\vec{x})} , \quad (2.7)$$

and

$$\mathcal{R}(\Omega) \hat{U}_k(\vec{x}) \mathcal{R}^\dagger(\Omega) = \Omega^\dagger(\vec{x}) \hat{U}_k(\vec{x}) \Omega(\vec{x} + \vec{e}_k) . \quad (2.8)$$

The vacuum state $|0\rangle$ is supposed to be gauge invariant

$$\mathcal{R}(\Omega)|0\rangle = |0\rangle ; \quad \Gamma^a(\vec{x})|0\rangle = 0 . \quad (2.9)$$

The second equation means that the electric flux equals to zero if there are no sources (Gauss law). It is easy to show that operator

$$P_0 = \int [d\Omega] \mathcal{R}(\Omega) \equiv \prod_{\vec{x}} \int d\Omega(\vec{x}) \exp \left\{ i \sum_a \omega^a(\vec{x}) \Gamma^a(\vec{x}) \right\} . \quad (2.10)$$

is the projection operator on the gauge invariant states.

Let H be a Hamiltonian (to be defined later). Then the partition function Z is

$$Z = \text{Tr} \left(\exp \left\{ -\frac{1}{T} H \right\} \right)_{colorless} ; \quad T \equiv \frac{1}{a_4 N_4} , \quad (2.11)$$

where T is the temperature and the trace is defined on some colorless space of states. By definition, the transfer matrix $\hat{V}$ is given by $\hat{V} = \exp\{-a_4 H\}$. Therefore,

$$Z = \text{Tr} \left(\hat{V}^{N_4} \right)_{colorless} \equiv \text{Tr} \left(\hat{V}^{N_4} P_0 \right) . \quad (2.12)$$

Let $|\Psi_k\rangle$ be eigenstates of the transfer matrix $\hat{V}$ with eigenvalues λ_k

$$\hat{V}|\Psi_k\rangle = \lambda_k |\Psi_k\rangle ; \quad \lambda_k \equiv e^{-a_4 E_k} , \quad (2.13)$$

and E_k are eigenvalues of the Hamiltonian. One can choose

$$E_0 < E_1 \leq E_2 \leq \dots . \quad (2.14)$$

Vacuum $|\text{vac}\rangle$ is the eigenstate with the lowest E_0 : $|\text{vac}\rangle \equiv |\Psi_0\rangle$, and it is supposed to be nondegenerate. The partition function Z can be represented as

$$Z = \text{Tr} \left(\hat{V}^{N_4} P_0 \right) \equiv \sum_{k \geq 0} \langle \Psi_k | \hat{V}^{N_4} P_0 | \Psi_k \rangle = \sum_{k \geq 0} \exp \left\{ -\frac{1}{T} E_k \right\} \cdot \langle \Psi_k | P_0 | \Psi_k \rangle . \quad (2.15)$$

Every eigenstate can be expanded as follows

$$|\Psi_k\rangle = \int [dU] \Psi_k(U) |U\rangle . \quad (2.16)$$

The set of the eigenstates $\{|\Psi_k\rangle\}$ is supposed to be complete

$$\sum_k \Psi_k^*(U') \Psi_k(U) = \delta(U; U') . \quad (2.17)$$

Then

$$\begin{aligned} Z &= \sum_{k \geq 0} \langle \Psi_k | \hat{V}^{N_4} P_0 | \Psi_k \rangle = \sum_{k \geq 0} \int [dU] [dU'] \Psi_k^*(U') \Psi_k(U) \cdot \langle U' | \hat{V}^{N_4} P_0 | U \rangle \\ &= \int [dU] \langle U | \hat{V}^{N_4} P_0 | U \rangle . \end{aligned} \quad (2.18)$$

Evidently, in the zero temperature limit the main contribution to the partition function Z comes from the vacuum eigenstate

$$Z = \langle \text{vac} | \exp \left\{ -\frac{1}{T} H \right\} | \text{vac} \rangle ; \quad T \equiv \frac{1}{a_4 N_4} \rightarrow 0 . \quad (2.19)$$

The partition function Z defined above presumes the time periodicity

$$U_\mu(\vec{x}; x_4 = L_4) = U_\mu(\vec{x}; x_4 = 0) . \quad (2.20)$$

By applying the gauge transformation $\Omega(\vec{x}; x_4)$

$$\Omega(\vec{x}; x_4) = \begin{cases} U_4^\dagger(\vec{x}; L_4 - a_4) ; & x_4 = L_4 - a_4 ; \\ \left(U_4(\vec{x}; L_4 - 2a_4) U_4(\vec{x}; L_4 - a_4) \right)^\dagger ; & x_4 = L_4 - 2a_4 ; \\ \dots \dots & \dots \dots \\ \left(U_4(\vec{x}; a_4) U_4(\vec{x}; 2a_4) \dots U_4(\vec{x}; L_4 - a_4) \right)^\dagger ; & x_4 = a_4 , \end{cases} \quad (2.21)$$

one obtains $U_4(\vec{x}; x_4) = 1$ at $x_4 \neq 0$. Let us define $\Lambda(\vec{x}) \equiv U_4(\vec{x}; 0)$, and

$$U_k^\Lambda(\vec{x}; 0) = \Lambda^\dagger(\vec{x}) U_k(\vec{x}; 0) \Lambda(\vec{x} + \hat{e}_k) . \quad (2.22)$$

Evidently, $\text{Tr } U_{P_s}^\Lambda(x_4) = \text{Tr } U_{P_s}(x_4)$. Defining functions W_E, W_M :

$$\begin{aligned} W_E(0) &= \frac{2Na}{g_t^2 a_4} \sum_{k; \vec{x}} \left[1 - \frac{1}{N} \text{ReTr} \left(U_k^\Lambda(\vec{x}; 0) U_k^\dagger(\vec{x}; a_4) \right) \right] ; & x_4 = 0 ; \\ W_E(x_4) &= \frac{2Na}{g_t^2 a_4} \sum_{k; \vec{x}} \left[1 - \frac{1}{N} \text{ReTr} \left(U_k(\vec{x}; x_4) U_k^\dagger(\vec{x}; x_4 + a_4) \right) \right] ; & x_4 \geq a_4 ; \\ W_M(x_4) &= \frac{2Na_4}{g_s^2 a} \sum_{P_s} \left(1 - \frac{1}{N} \text{ReTr} U_{P_s}(x_4) \right) , \end{aligned} \quad (2.23)$$

one obtains

$$S(U) = \sum_{x_4=0}^{L_4-a_4} \left\{ W_E(x_4) + \frac{1}{2} [W_M(x_4) + W_M(x_4 + a_4)] \right\} . \quad (2.24)$$

Let us introduce the (pure gauge) transfer matrix $\hat{V}_G$ with matrix elements

$$\langle U(x_4 + a_4) | \hat{V}_G | U(x_4) \rangle = C_0^{-1} \cdot \exp \left\{ -W_E(x_4) - \frac{1}{2} [W_M(x_4) + W_M(x_4 + a_4)] \right\} , \quad (2.25)$$

where C_0 is some constant. Then

$$Z = \int [d\Lambda] \int \prod_{x_4=0}^{L_4-a_4} [dU(x_4)] \exp \left\{ - \sum_{x_4} [W_E(x_4) + W_M(x_4)] \right\} = C_0^{N_4} \cdot \text{Tr} \left(\hat{V}_G^{N_4} P_0 \right) , \quad (2.26)$$

where

$$[d\Lambda] = \prod_{\vec{x}} d\Lambda(\vec{x}) ; \quad [dU(x_4)] = \prod_{\vec{x}} \prod_k dU_k(\vec{x}; x_4) , \quad (2.27)$$

and P_0 is the projection operator on the colorless state defined above.

Let us note that in the finite volume one obtains the standard Wilson action only in the case of the periodic boundary condition, i.e. when $W_M(L_4) = W_M(0)$.

3 Disorder parameter $\langle \hat{\mu} \rangle$

For every spacelike link $l_s = (\vec{x}; k)$ let us define matrices

$$\overline{\Omega}_{l_s} \equiv \exp\{i\vec{b}_{l_s}\vec{T}\} \in SU(N) , \quad (3.1)$$

where $\vec{b}_{l_s}$ are any parameters (not necessarily connected with monopoles). Following the Pisa approach one can define the disorder operator

$$\hat{\mu}(\vec{b}) = \exp\left\{i \sum_{l_s} \vec{b}_{l_s} \vec{E}_{l_s}\right\} . \quad (3.2)$$

Then the disorder parameter $\langle \hat{\mu} \rangle$ is given by

$$\langle \hat{\mu} \rangle = \frac{1}{Z} Z(\overline{\Omega}) ; \quad Z(\overline{\Omega}) = \text{Tr} \left(\prod_{l_s} R_{l_s}(\overline{\Omega}_{l_s}) \hat{V}_G^{N_4} P_0 \right) , \quad (3.3)$$

where operators $R_{l_s}(\overline{\Omega}_{l_s})$, $\hat{V}_G$ and P_0 have been defined in the previous section. Repeating the procedure of the previous section, one obtains in the temporal gauge (tg) :

$$Z(\overline{\Omega}) = \int [dU]_{tg} e^{-S(U; \overline{\Omega})} ; \quad [dU]_{tg} \equiv \prod_{\vec{x}} dU_4(\vec{x}, 0) \prod_{x_4=0}^{L_4-a} \prod_{\vec{x}k} dU_k(\vec{x}; x_4) , \quad (3.4)$$

where

$$S(U; \overline{\Omega}) = S_W(U) + \delta S_E(U; \overline{\Omega}) + \delta S_M(U; \overline{\Omega}) ;$$

$$\delta S_E = \frac{2}{g^2} \sum_{k; \vec{x}} \text{ReTr} \left[(1 - \overline{\Omega}_{\vec{x}; k}) \cdot U_k(\vec{x}; L_4 - a) U_k^\dagger(\vec{x}; 0) \right] ; \quad (3.5)$$

$$\delta S_M = \frac{1}{g^2} \sum_{i < k} \sum_{\vec{x}} \text{ReTr} \left[U_{\vec{x}; i} U_{\vec{x}+\hat{i}; k} U_{\vec{x}+\hat{k}; i}^\dagger U_{\vec{x}; k}^\dagger - U'_{\vec{x}; i} \cdot U'_{\vec{x}+\hat{i}; k} \cdot U'_{\vec{x}+\hat{k}; i}{}^\dagger \cdot U'_{\vec{x}; k}{}^\dagger \right] (x_4 = 0) .$$

and $U'_{l_s} = \overline{\Omega}_{l_s}^\dagger U_{l_s}$. To get rid of the temporal gauge one can perform successively the change of variables $U \rightarrow U''$:

$$U_k(\vec{x}; x_4) = U_4^\dagger(\vec{x}; x_4) U_k''(\vec{x}; x_4) U_4(\vec{x} + \hat{k}; x_4) ; \quad (3.6)$$

$$U_4(\vec{x}; x_4 - a) = U_4''(\vec{x}; x_4 - a) U_4(\vec{x}; x_4) \quad (3.7)$$

at $x_4 = a$, then at $x_4 = 2a$, ... , then at $x_4 = L_4 - a$. Finally, one arrives at

$$\langle \hat{\mu}(\overline{\Omega}) \rangle = \frac{1}{Z} \int [dU] e^{-\tilde{S}(U; \overline{\Omega})} ; \quad \tilde{S}(U; \overline{\Omega}) = S_W(U) + \delta S_E(U; \overline{\Omega}) ; \quad (3.8)$$

$$\delta S_E(U; \overline{\Omega}) = \frac{2}{g^2} \sum_{k; \vec{x}} \text{ReTr} \left[(1 - \overline{\Omega}_{\vec{x}; k}) \cdot U_{\vec{x}; k}(0) U_{\vec{x}+\hat{k}; 4}(0) U_{\vec{x}; k}^\dagger(a) U_{\vec{x}; 4}^\dagger(0) \right] . \quad (3.9)$$

Equations (3.8) and (3.9) define the desired – functional – representation of the order parameter $\langle \hat{\mu} \rangle$.

4 Summary

Finally, let us summarize. This paper is dedicated to the discussion of some formal aspects of the functional integral representation of the disorder parameter in lattice gauge theories. No special choice of the fields $\vec{b}_{l_s}$ has been used.

The main results are the following.

- 1) The functional integral representation of the disorder parameter $\langle \hat{\mu} \rangle$ has been derived in the canonical quantization approach. This derivation resolves the problem – discussed in paper [5] – of the (unambiguous) choice of the modified action $\tilde{S}(U; \overline{\Omega})$ on the lattice.

The representation obtained for the disorder parameter $\langle \hat{\mu} \rangle$ is given in eq.'s (3.8) and (3.9) and coincides with that used, say, in paper [10].

- 2) As a byproduct, this derivation establishes the boundary conditions along the imaginary time direction : they have to be periodic.

The choice of, say, C^* -periodic boundary conditions (see, e.g., [10]) is not compatible with the canonical quantization approach.

- 3) Also, this derivation resolves the problem of the choice of the gauge in the functional integral representation for $\langle \hat{\mu} \rangle$: no gauge fixing is needed.

This is in agreement with the numerical results of the paper [10] where the so called RAP (i.e. no gauge fixing) has been also used.

Acknowledgments

This work has been supported by the RFBR grant 05-02-16306, DFG–RFBR grant 06-02-04014 and Heisenberg–Landau program.

References

- [1] L.P. Kadanoff and H. Ceva, Phys. Rev. **B3** (1971) 3918.
- [2] G. 't Hooft, in *High Energy Physics*, Proceedings of the EPS International Conference, Palermo 1975, ed. A. Zichichi, Editrice Compositori, Bologna 1976.
- [3] S. Mandelstam, Phys. Rep. **23** (1976) 245.
- [4] H.B. Nielsen and P. Olesen, Nucl. Phys. **B61** (1973) 61.
- [5] L. Del Debbio, A. Di Giacomo and G. Paffuti, Phys. Lett. **B349** (1995) 513.
- [6] A. Di Giacomo and G. Paffuti, Phys. Rev. **D56** (1997) 5816.
- [7] L. Del Debbio, A. Di Giacomo, G. Paffuti and P. Pieri, Phys. Lett. **B355** (1995) 255.
- [8] A. Di Giacomo, B. Lucini, L. Montesi and G. Paffuti, Phys. Rev. **D61** (2000) 034503.
- [9] A. Di Giacomo, B. Lucini, L. Montesi and G. Paffuti, Phys. Rev. **D61** (2000) 034504.
- [10] J.M. Carmona, M. D'Elia, A. Di Giacomo, B. Lucini and G. Paffuti, Phys. Rev. **D64** (2001) 114507.
- [11] A. Di Giacomo, hep-lat/0206018 (2002).
- [12] M. D'Elia, A. Di Giacomo, B. Lucini, G. Paffuti and C. Pica, Phys. Rev. **D71** (2005) 114502.
- [13] A. Barresi, G. Burgio, M. D'Elia and M. Mueller-Preussker, Phys. Lett. **B599** (2004) 278.
- [14] G. Burgio, M. Fuhrmann, W. Kerler and M. Mueller-Preussker, hep-lat/0610097 (2006).
- [15] M. N. Chernodub, M. I. Polikarpov and A. I. Veselov, Phys. Lett. **B399** (1997) 267; Nucl. Phys. Proc. Suppl. **49** (1996) 307;
- [16] V. A. Belavin, M. N. Chernodub and M. I. Polikarpov, JETP Lett. **83** (2006) 308.
- [17] J. Frohlich and P. A. Marchetti, Commun. Math. Phys. **112** (1987) 343.